



## MEET KELLY



42-year-old female



Digital artist; spends most of the day working on a tablet or laptop



### Medical History

- **No significant** ocular history
- Has self-medicated with **multiple OTC drops over the last year** but isn't sure which ones; found them **ineffective**
- Takes a **daily antihistamine**



### Chief Complaints

- Increasing **difficulty with near vision**; patient **believes she requires glasses**
- Symptoms of **eye fatigue, burning eyes, and fluctuating vision**, especially late in the day
- Reports a **gritty and scratchy feeling** in her eyes. Visual symptoms and dry, irritated eyes **affect her ability to perform her work**



### Initial Observations

- Mild eye **redness**
- **Blinks to improve vision** during visual acuity testing

### INDICATION

MIEBO™ (perfluorohexyloctane ophthalmic solution) is a semifluorinated alkane indicated for the treatment of the signs and symptoms of dry eye disease.

### IMPORTANT SAFETY INFORMATION

- MIEBO should not be administered while wearing contact lenses. Contact lenses should be removed before use and for at least 30 minutes after administration of MIEBO

**Please see additional Important Safety Information on next page.**

**Click [here](#) for full Prescribing Information for MIEBO.**

**Miebo**<sup>™</sup>  
(perfluorohexyloctane  
ophthalmic solution)

# NEXT STEPS



Based on Kelly's history and presentation, **what assessments would you perform?**



**What results would you expect to find?**



What would be **your proposed treatment plan** to address Kelly's symptoms?



Discover how MIEBO could provide patients like Kelly with **relief from the signs and symptoms** of dry eye disease.

**Learn more at [MIEBO-ECP.COM](https://www.miebo-ecp.com).**

## IMPORTANT SAFETY INFORMATION (CONTINUED)

- Instruct patients to instill one drop of MIEBO into each eye four times daily
- The safety and efficacy in pediatric patients below the age of 18 have not been established
- The most common ocular adverse reaction was blurred vision (1% to 3% of patients reported blurred vision and conjunctival redness)

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](https://www.fda.gov/medwatch) or call 1-800-FDA-1088.

**Please see additional Important Safety Information on previous page.**

**Click [here](#) for full Prescribing Information for MIEBO.**

**BAUSCH + LOMB**

©2023 Bausch + Lomb  
MBO.0406.USA.23

**Miebo**<sup>™</sup>  
(perfluorohexyloctane  
ophthalmic solution)