

MEET MALIK

- 36-year-old male
- Web developer and video game enthusiast



Medical History

 Occasional eye discomfort and fluctuating vision led to appointment to update glasses prescription



Chief Complaints

- Blurry vision continues to be bothersome; temporarily improves with blinking
- Symptoms are beginning to affect work



Initial Observations

• Tired-looking eyes, signs of irritation

INDICATION

MIEBO™ (perfluorohexyloctane ophthalmic solution) is a semifluorinated alkane indicated for the treatment of the signs and symptoms of dry eye disease.

IMPORTANT SAFETY INFORMATION

 MIEBO should not be administered while wearing contact lenses. Contact lenses should be removed before use and for at least 30 minutes after administration of MIEBO

Please see additional Important Safety Information on next page. Click <u>here</u> for full Prescribing Information for MIEBO.





NEXT STEPS



Based on Malik's history and presentation, what assessments would you perform?



What results would you expect to find?



What would be **your proposed treatment plan** to address Malik's symptoms?





Discover how MIEBO could provide patients like Malik with relief from the signs and symptoms of dry eye disease.

Learn more at MIEBO-ECP.COM.



IMPORTANT SAFETY INFORMATION (CONTINUED)

- Instruct patients to instill one drop of MIEBO into each eye four times daily
- The safety and efficacy in pediatric patients below the age of 18 have not been established
- The most common ocular adverse reaction was blurred vision (1% to 3% of patients reported blurred vision and conjunctival redness)

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see additional Important Safety Information on previous page. Click <u>here</u> for full Prescribing Information for MIEBO.





