

The first and only Rx eye drop for DED that directly targets evaporation¹

Product Information

PRODUCT TRADE: MIEBO

DELIVERY: Topical ophthalmic solution

ACTIVE INGREDIENT:

Perfluorohexyloctane 100%

INACTIVE INGREDIENTS: None

PRESERVATIVE: None


HOW SUPPLIED: Polypropylene bottles with dropper tips and screw caps in the following size: 3-mL fill in a 5-mL bottle

INDICATION AND USAGE: MIEBO is indicated for the treatment of the signs and symptoms of dry eye disease (DED)

FILL SIZE: 3-mL in 5-mL bottle

NO A/B GENERIC EQUIVALENT AVAILABLE

MIEBO DOSING: Instill one drop of MIEBO four times daily into affected eye(s).



One drop per eye, QID dosing¹



QID, 4 times daily.

1 bottle = 1 month

The MIEBO drop is small (11 µL)*

~270 drops, or 1-month supply

Convenient, preservative-free, multidose bottle^{1,2}

*Aqueous formulations have a typical drop size of ~30–50 µL.²

MIEBO contains only ONE ingredient¹



Water free

Preservative free

Steroid free

Vehicle free

What patients can expect

- The small drop size (11 µL) means **MIEBO may feel different** from formulations containing water²
- There may be **no ocular sensation or blink reflex** upon instillation.³

See Section 2.2 of the Prescribing Information for administration instructions.

INDICATION

MIEBO® (perfluorohexyloctane ophthalmic solution) is indicated for the treatment of the signs and symptoms of dry eye disease.

IMPORTANT SAFETY INFORMATION

- MIEBO should not be administered while wearing contact lenses. Contact lenses should be removed before use and for at least 30 minutes after administration of MIEBO

Please see additional Important Safety Information on the back and [click here](#) for full Prescribing Information.

Order as usual: Each unit (1 bottle) of MIEBO contains a 1-month supply of drops

Packaging specifications

SELLING UNIT

| Unit | NDC | Dimensions | Weight |
|------|---------------|------------------------------|---------|
| 1 | 24208-0377-05 | 3.5 in x 1.375 in x 1.375 in | 0.04 lb |

SHIPPING CASE

| Pack | Dimensions | Weight |
|------|---------------------------|--------|
| 48 | 8.875 in x 6 in x 7.75 in | 2.3 lb |

BAR CODING:

3-mL fill in a 5-mL container bar-coded with NDC# 24208-0377-05

STORAGE CONDITIONS:

Store MIEBO at 15 °C to 25 °C (59 °F to 77 °F). After opening, MIEBO can be used until the expiration date on the bottle

EXPIRATION: 24 months

INNER PACK: 1

Wholesaler order numbers

| Fill | NDC | Package size | Cardinal | McKesson | Amerisource Bergen |
|-------------------------------|---------------|------------------------------|----------|----------|--------------------|
| 3-mL fill in a 5-mL container | 24208-0377-05 | 3.5 in x 1.375 in x 1.375 in | 5858964 | 2841351 | 10281423 |

Bausch + Lomb customer service department

Phone orders: 1-800-321-4576, Option 1

Fax orders: 1-908-927-1926

Email orders: Ophthapharma@bausch.com



For more information about MIEBO, including our patient savings programs, please visit www.miebo-ecp.com.

IMPORTANT SAFETY INFORMATION (CONTINUED)

- Instruct patients to instill one drop of MIEBO into each eye four times daily
- The safety and efficacy in pediatric patients below the age of 18 have not been established
- The most common ocular adverse reaction was blurred vision (1% to 3% of patients reported blurred vision and conjunctival redness)

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see additional Important Safety Information on the front and [click here](#) for full Prescribing Information.

References: 1. MIEBO. Prescribing Information. Bausch & Lomb, Inc. 2. Sheppard JD, Nichols KK. Dry eye disease associated with meibomian gland dysfunction: focus on tear film characteristics and the therapeutic landscape. *Ophthalmol Ther.* 2023;12(3):1397-1418. doi:10.1007/s40123-023-00669-1 3. Schmidl D, Bata AM, Szegedi S, et al. Influence of perfluorohexyloctane eye drops on tear film thickness in patients with mild to moderate dry eye disease: a randomized controlled clinical trial. *J Ocul Pharmacol Ther.* 2020;36(3):154-161. doi:10.1089/jop.2019.0092

BAUSCH + LOMB

©2024 Bausch + Lomb. MBO.0396.USA.24

Miebo[®]
(perfluorohexyloctane
ophthalmic solution)